

Safe and Tight Linear Estimators for Global Optimization

Glencora Borradaile and Pascal Van Hentenryck

Department of Computer Science
Brown University
Providence, Rhode Island 02912

CS-03-11
June 2003

Safe and Tight Linear Estimators for Global Optimization

Glencora Borradaile and Pascal Van Hentenryck

Department of Computer Science,
Brown University,
Box 1910,
Providence, RI, 02912
{glencora, pvh}@cs.brown.edu

Abstract. Global optimization problems are often approached by branch and bound algorithms which use linear relaxations of the nonlinear constraints computed from the current variable bounds. This paper studies how to derive safe linear relaxations to account for numerical errors arising when computing the linear coefficients. It first proposes two classes of safe linear estimators for univariate functions. Class-1 estimators generalize previously suggested estimators from quadratic to arbitrary functions, while class-2 estimators are novel. Class-2 estimators are shown to be tighter theoretically (in a certain sense) and almost always tighter numerically. The paper then generalizes these results to multivariate functions. It shows how to derive estimators for multivariate functions by combining univariate estimators derived for each variable independently. Moreover, it shows that the combination of tight class-1 safe univariate estimators gives a tight class-1 safe multivariate estimators.

1 Introduction

Global optimization problems arise naturally in many application areas, including chemical and electrical engineering, biology, economics, and robotics to name only a few. They consist of finding all solutions or the global optima to nonlinear programming problems. These problems are inherently difficult computationally (i.e., they are PSPACE-hard) and may also be challenging numerically. In addition, there has been considerable interest in recent years to produce rigorous or reliable results, i.e., to make sure that the exact solutions are enclosed in the results of the algorithms.

Global optimization problems are often approached by branch and bound algorithms which use linear relaxations of the nonlinear constraints computed from the current bounds for the variables at each node of the search tree (e.g., [3, 5, 6, 11, 12, 15, 16]). The linear relaxation can be used to obtain a lower bound on the objective function (in minimization problems) and/or to update the variable bounds. This approach can also be combined with constraint satisfaction techniques for global optimization which are also effective in reducing the variable bounds (e.g., [1, 2, 7, 13, 14]).

The linear relaxation is generally obtained by linearizing nonlinear terms independently, giving what is often called linear over- and under-estimators. When rigorous and reliable results are desired, it is critical to generate a safe linear relaxation which overapproximates the solution to the nonlinear problem at hand (e.g., [9, 10]). Indeed, the coefficients in the linear constraints are generally given by real functions which are subject to rounding errors when evaluated. As a consequence, the resulting linear relaxation may not be safe. Moreover, naive approaches (e.g., upward rounding for overestimators) are not safe either in general. Once a safe linear relaxation is available, it can be solved exactly or safe bounds on the objective function can be obtained using duality as in [10, 4] for instance. Experimental results (e.g., [9]) have shown that both of these two corrections are critical in practice, even on simple problems, to find all solutions to nonlinear polynomial systems.

This paper focuses on obtaining safe linear relaxations for global optimization problems and contains two main contributions:

1. The paper presents two classes of safe estimators of univariate functions. The first class of estimators generalizes the results of [9] from quadratic to arbitrary functions, while the second class is entirely novel. Theoretical tightness results are given for both classes, giving the relative strengths of the presented estimators. In particular, the results show that class-2 estimators (when they apply) are theoretically tighter than class-1 estimators (in a certain sense to be defined). Moreover, the numerical results indicate that class-2 estimators are almost always tighter in practice in our experiments.
2. The paper then generalizes the univariate results to multivariate functions. It shows how to derive estimators for multivariate functions by combining univariate estimators derived for each variable independently. Moreover, it proves that the combination of tight class-1 safe univariate estimators is a tight class-1 safe multivariate estimator.

As a consequence, these results provide a systematic, comprehensive, and elegant framework to derive safe linear estimators for global optimization problems. In conjunction with the safe bounds on the linear relaxations derived in [10, 4], they provide the theoretical foundation for rigorous results of branch and bound approaches to global optimization based on linear programming.

The rest of this paper is organized as follows: Section 2 defines the concept of safe estimators and the problems arising in deriving them. Section 3 derives the two classes of linear estimators for univariate functions. Section 4 presents the theoretical and numerical tightness results. Section 5 presents the multivariate results.

2 Definitions and Problem Statement

This section defines the concepts of linear estimators and safe linear overestimators, as well as the problem tackled in this paper. For simplicity, the definitions are given for univariate functions only, but they generalize naturally

to multivariate functions. We only consider overestimators, since the treatment for underestimators is similar. A linear overestimator is a linear function which provides an upper bound to a univariate function over an interval.

Definition 1 (Linear Overestimators). *Let g be a univariate function $\mathbb{R} \rightarrow \mathbb{R}$. A linear overestimator of g over the interval $[\underline{x}, \overline{x}]$ is a linear function $mx + b$ satisfying*

$$\forall x \in [\underline{x}, \overline{x}] : mx + b \geq g(x).$$

In general, given a univariate function g , linear overestimators are obtained through tangent or secant lines. These are implicitly specified by two functions $f_m(\underline{x}, \overline{x}, g)$ and $f_b(\underline{x}, \overline{x}, g)$ computing the slope m and the intercept b of the estimator.¹ For instance, the secant line for the function x^n (n even) is the linear overestimator $mx + b$ over $[\underline{x}, \overline{x}]$ specified by

$$m = \frac{\overline{x}^n - \underline{x}^n}{\overline{x} - \underline{x}} \quad b = \frac{\overline{x}\overline{x}^n - \underline{x}\underline{x}^n}{\overline{x} - \underline{x}}$$

Unfortunately, given an underlying floating-point system $\mathcal{F}$ and a representation of f_m and f_b , the computation of these functions is subject to rounding errors and will produce the approximations $\tilde{m}$ and $\tilde{b}$. However, the linear function $\tilde{m}x + \tilde{b}$ is not guaranteed to be a linear estimator of g in $[\underline{x}, \overline{x}]$.

The main issue addressed in this paper is how to compute safe linear overestimators, i.e., linear overestimators $m^*x + b^*$ where m^* and b^* are floating-point numbers in the underlying floating-point system $\mathcal{F}$.

Definition 2 (Safe Linear Overestimators). *Let g be a univariate function $\mathbb{R} \rightarrow \mathbb{R}$. A safe linear overestimator of g over interval $[\underline{x}, \overline{x}]$ is a linear overestimator $m^*x + b^*$ for g over $[\underline{x}, \overline{x}]$ where $m^*, b^* \in \mathcal{F}$.*

Notations In the following, we often abuse notation and use m and b to represent the functions f_m and f_b . The critical point to remember is that m and b cannot be computed exactly and may involve significant rounding errors. Also, given an expression e , we use $\lfloor e \rfloor$ and $\lceil e \rceil$ to denote the most precise lower and upper approximation of e at our disposal given $\mathcal{F}$ and the representation of e .²

The Problem At first sight, it may seem that the problem of finding a safe linear overestimator $m^*x + b^*$ is trivial: simply choose the function $\lceil m \rceil x + \lceil b \rceil$, i.e., choose $m^* = \lceil m \rceil$ and $b^* = \lceil b \rceil$. Unfortunately, as shown in Figure 1, this is not correct. The figure shows g and its linear overestimator $mx + b$ over $[\underline{x}, \overline{x}]$. The estimator is correct in the $\mathbb{R}^+$ region, but not in the $\mathbb{R}^-$ region where the slope $\lceil m \rceil$ is too strong. Similarly, $\lfloor m \rfloor x + \lceil b \rceil$ is not a safe overestimator because its slope is too weak in the $\mathbb{R}^+$ region. The value b^* must be chosen carefully even when $m^* = \lceil m \rceil$ or when $m^* = \lfloor m \rfloor$. The figure shows such a choice of b^* .

¹ More precisely, they are specified by a representation (e.g., the text) of these two functions.

² Note the safety results presented in this paper hold even if the approximations are not the most precise.




Fig. 1. Why $\lceil m \rceil x + \lceil b \rceil$ is not a Safe Linear Overestimator.

Tightness In addition to safety, one is generally interested in linear overestimators which are as tight as possible given $\mathcal{F}$ and the representation of f_m and f_b .

Definition 3 (Error of Linear Overestimators). Let g be a univariate function $\mathbb{R} \rightarrow \mathbb{R}$. The error of a linear overestimator $mx + b$ for g over $[\underline{x}, \bar{x}]$ is given by

$$\int_{\underline{x}}^{\bar{x}} mx + b - g(x) dx.$$

Definition 4 (Tightness of Linear Overestimators). Let g be a univariate function $\mathbb{R} \rightarrow \mathbb{R}$. Let l_1 and l_2 be two linear overestimators of g over $[\underline{x}, \bar{x}]$. l_1 is tighter than l_2 wrt g and $[\underline{x}, \bar{x}]$ if l_1 has a smaller error than l_2 ,

3 Safe Linear Overestimators for Univariate Functions

This section describes two classes of safe overestimators. These estimators are derived from the linear overestimator $mx + b$. In other words, the goal is to find m^* and b^* in $\mathcal{F}$ such that

$$m^*x + b^* \geq mx + b, \quad \forall x \in [\underline{x}, \bar{x}]. \quad (1)$$

As mentioned, the first class generalizes the results of Michel et al. [9] who gave safe estimators for x^2 . The second class is entirely new and enjoys some nice theoretical and numerical properties.




Fig. 2. A Safe Overestimator when $\underline{x} \geq 0$.

3.1 A First Class of Safe Overestimators

To obtain a safe linear overestimator $m^*x + b^*$ from $mx + b$, there are only two reasonable choices for m^* : $\lfloor m \rfloor$ and $\lceil m \rceil$. Other choices would necessarily be less tight. We derive the safe overestimators for $\lceil m \rceil$ only, the derivation for $\lfloor m \rfloor$ being similar. The problem thus reduces to finding b^* such that

$$\lceil m \rceil x + b^* \geq mx + b. \quad (2)$$

Since $\lceil m \rceil \geq m$, it is sufficient to satisfy (2) at $x = \underline{x}$, which implies $b^* \geq b - (\lceil m \rceil - m)\underline{x}$. The overestimator can now be derived by a case analysis on the sign of $\underline{x}$. If $\underline{x} \leq 0$, we have

$$b - (\lceil m \rceil - m)\underline{x} \leq b - (\lceil m \rceil - \lfloor m \rfloor)\underline{x} = b - \mathbf{err}(m)\underline{x}$$

where $\mathbf{err}(m) = \lceil m \rceil - \lfloor m \rfloor$. Therefore, choosing $b^* = \lceil b - \mathbf{err}(m)\underline{x} \rceil$ satisfies (2). If $\underline{x} \geq 0$, it is sufficient to choose $b^* = \lceil b \rceil$, as shown in Figure 2. The following theorem summarizes the results.

Theorem 1 (Safe Linear Overestimators for Univariate Functions, Class 1). *Let g be a univariate function g and let $mx + b$ be a linear overestimator for g in $[\underline{x}, \bar{x}]$. We have that*

$$mx + b \leq \begin{cases} \lfloor m \rfloor x + \lceil b + \mathbf{err}(m)\bar{x} \rceil & \text{if } \bar{x} \geq 0 \\ \lfloor m \rfloor x + \lceil b \rceil & \text{if } \bar{x} \leq 0 \\ \lceil m \rceil x + \lceil b - \mathbf{err}(m)\underline{x} \rceil & \text{if } \underline{x} \leq 0 \\ \lceil m \rceil x + \lceil b \rceil & \text{if } \underline{x} \geq 0. \end{cases}$$




Fig. 3. Choosing $b^* = \lceil g(\underline{x}) - \lfloor m \rfloor \underline{x} \rceil$ which is almost always tighter than $\lceil m \rceil x + \lceil b \rceil$.

As a consequence, the four right-hand sides are safe linear overestimators for g in $[\underline{x}, \bar{x}]$ under the specified conditions.

Proof. We give the proofs for the first two cases. The proofs are similar for the symmetric cases. In the first case, we have

$$\begin{aligned}
 & \lfloor m \rfloor x + \lceil b + \text{err}(m) \bar{x} \rceil \\
 &= \lfloor m \rfloor (\bar{x} - s) + \lceil b + \text{err}(m) \bar{x} \rceil && \text{letting } x = \bar{x} - s \text{ with } s \geq 0 \\
 &\geq \lfloor m \rfloor (\bar{x} - s) + b + \text{err}(m) \bar{x} \\
 &= \lfloor m \rfloor (\bar{x} - s) + b + (\lceil m \rceil - \lfloor m \rfloor) \bar{x} \\
 &= \lceil m \rceil \bar{x} - \lfloor m \rfloor s + b \\
 &\geq m \bar{x} - \lfloor m \rfloor s + b && \lceil m \rceil \bar{x} \geq m \bar{x} \text{ since } \bar{x} \geq 0 \\
 &\geq m \bar{x} - ms + b && \lfloor m \rfloor s \leq ms \text{ since } s \geq 0 \\
 &= mx + b
 \end{aligned}$$

In the second case, since $x \leq \bar{x} \leq 0$, we have that $mx \leq \lfloor m \rfloor x$ and hence $mx + b \leq \lfloor m \rfloor x + \lceil b \rceil$. $\square$

3.2 A Second Class of Safe Overestimators

In general, the overestimators used in global optimization are either secant lines of g (as in Figure 1) or tangent lines to g at $\underline{x}$ or $\bar{x}$ (see, for instance, [5]). As a consequence, we have that $g(\underline{x}) = m \underline{x} + b$ and/or $g(\bar{x}) = m \bar{x} + b$. In these circumstances, it is possible to find a b^* satisfying (2) which does not depend on

the sign of $\underline{x}$. Assume that $g(\underline{x}) = m\underline{x} + b$. Since b^* must satisfy $\lceil m \rceil \underline{x} + b^* \geq m\underline{x} + b = g(\underline{x})$, it follows that $b^* \geq g(\underline{x}) - \lceil m \rceil \underline{x}$. Choosing $b^* = \lceil g(\underline{x}) - \lceil m \rceil \underline{x} \rceil$ also satisfies (2). This choice for b^* enjoys some nice theoretical and experimental properties as detailed in Section 4. Figure 3 illustrates this situation.

Theorem 2 (Safe Linear Overestimators of Univariate Functions, Class 2). *Let g be a univariate function g and let $m\underline{x} + b$ be a linear overestimator for g in $[\underline{x}, \overline{x}]$. We have that*

$$m\underline{x} + b \leq \begin{cases} \lfloor m \rfloor \underline{x} + \lceil g(\overline{x}) - \lfloor m \rfloor \overline{x} \rceil & \text{if } g(\overline{x}) = m\overline{x} + b \\ \lceil m \rceil \underline{x} + \lceil g(\underline{x}) - \lceil m \rceil \underline{x} \rceil & \text{if } g(\underline{x}) = m\underline{x} + b \end{cases}$$

As a consequence, the right-hand sides are safe linear overestimators for g in $[\underline{x}, \overline{x}]$ under the specified conditions.

Proof. We give the proof for the first case. The proof is similar for the symmetric case. We have

$$\begin{aligned} & \lfloor m \rfloor \underline{x} + \lceil g(\overline{x}) - \lfloor m \rfloor \overline{x} \rceil \\ &= \lfloor m \rfloor (\overline{x} - s) + \lceil g(\overline{x}) - \lfloor m \rfloor \overline{x} \rceil && \text{letting } \underline{x} = \overline{x} - s \text{ with } s \geq 0 \\ &\geq \lfloor m \rfloor (\overline{x} - s) + g(\overline{x}) - \lfloor m \rfloor \overline{x} \\ &= m\overline{x} + b - \lfloor m \rfloor s && \text{since } g(\overline{x}) = m\overline{x} + b \\ &\geq m\overline{x} + b - ms && \lfloor m \rfloor s \leq ms \text{ since } s \geq 0 \\ &= m\underline{x} + b. \end{aligned}$$

□

4 Tightness of Safe Linear Overestimators

Theorems 1 and 2 provide us with six safe overestimators of a univariate function. Several of the conditions for these estimators are not mutually exclusive and it is natural to study their relative tightness. Of course, it is possible to use all applicable safe estimators in the linear relaxation, but this may be undesirable for numerical and efficiency reasons. This section presents theoretical and experimental results on the tightness of the estimators.

4.1 Theoretical Results on Class 1 Estimators

This section studies the tightness of class-1 estimators. We first compare the estimators $\lfloor m \rfloor \underline{x} + \lceil b + \mathbf{err}(m)\overline{x} \rceil$ and $\lceil m \rceil \underline{x} + \lceil b - \mathbf{err}(m)\underline{x} \rceil$ which are both applicable when $0 \in [\underline{x}, \overline{x}]$. The result shows which estimators to choose according to the magnitude of $\underline{x}$ and $\overline{x}$.

Theorem 3 (Tightness of Class 1 Safe Linear Overestimators when $0 \in [\underline{x}, \overline{x}]$). *Let g be a univariate function g , $m\underline{x} + b$ be a linear overestimator for g in*




Fig. 4. Finding the optimal floating point representation. In this case $\lfloor m \rfloor x + b_2^*$ is a tighter estimator than $\lfloor m \rfloor x + b_1^*$.

$[\underline{x}, \overline{x}]$, and $\underline{x} < 0$ and $\overline{x} > 0$. The safe linear overestimator $\lfloor m \rfloor x + \lceil b + \text{err}(m)\overline{x} \rceil$ is tighter than the safe linear overestimator $\lfloor m \rfloor x + \lceil b - \text{err}(m)\underline{x} \rceil$ if $|\overline{x}| < |\underline{x}|$. Similarly, the safe linear overestimator $\lfloor m \rfloor x + \lceil b - \text{err}(m)\underline{x} \rceil$ is tighter than the safe linear overestimator $\lfloor m \rfloor x + \lceil b + \text{err}(m)\overline{x} \rceil$ if $|\underline{x}| < |\overline{x}|$.

Proof. To compare the estimators $\lfloor m \rfloor x + \lceil b + \text{err}(m)\overline{x} \rceil$ and $\lfloor m \rfloor x + \lceil b - \text{err}(m)\underline{x} \rceil$, we compare the relative tightness of the slightly tighter estimators $\lfloor m \rfloor x + b + \text{err}(m)\overline{x}$ and $\lfloor m \rfloor x + b - \text{err}(m)\underline{x}$. Their tightness is easier to determine and approximates well the actual relative tightness, since the rounding errors in computing $\lceil b + \text{err}(m)\overline{x} \rceil$ and $\lceil b - \text{err}(m)\underline{x} \rceil$ are comparable. First consider the error of $\lfloor m \rfloor x + b + \text{err}(m)\overline{x}$:

$$\begin{aligned}
 E_1 &= \int_{\underline{x}}^{\overline{x}} \lfloor m \rfloor x + b + \text{err}(m)\overline{x} - g(x) dx \\
 &= \int_{\underline{x}}^{\overline{x}} \lfloor m \rfloor x + b + \text{err}(m)\overline{x} - (mx + b) + mx + b - g(x) dx \\
 &= \int_{\underline{x}}^{\overline{x}} (\lfloor m \rfloor - m)x + \text{err}(m)\overline{x} dx + \underbrace{\int_{\underline{x}}^{\overline{x}} mx + b - g(x) dx}_E \\
 &= (\overline{x} - \underline{x}) \left[\overline{x} \left(\lfloor m \rfloor - \frac{1}{2} \lfloor m \rfloor - \frac{1}{2} m \right) + \underline{x} \left(\frac{1}{2} \lfloor m \rfloor - \frac{1}{2} m \right) \right] + E
 \end{aligned}$$

The error of $\lceil m \rceil x + b - \mathbf{err}(m)\underline{x}$ is similarly given by:

$$\begin{aligned} E_2 &= \int_{\underline{x}}^{\overline{x}} \lceil m \rceil x + b - \mathbf{err}(m)\underline{x} - g(x) dx \\ &= (\overline{x} - \underline{x}) \left[\overline{x} \left(\frac{1}{2} \lceil m \rceil - \frac{1}{2} m \right) + \underline{x} \left(\lfloor m \rfloor - \frac{1}{2} \lceil m \rceil - \frac{1}{2} m \right) \right] + E \end{aligned}$$

Estimator $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$ is tighter than estimator $\lceil m \rceil x + b - \mathbf{err}(m)\underline{x}$ when $E_1 < E_2$, i.e., when $|\overline{x}| < |\underline{x}|$. Similarly, estimator $\lceil m \rceil x + b - \mathbf{err}(m)\underline{x}$ is tighter than estimator $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$ when $|\underline{x}| < |\overline{x}|$. $\square$

When $\underline{x} \geq 0$, it is interesting to compare estimators $\lfloor m \rfloor x + \lceil b + \mathbf{err}(m)\overline{x} \rceil$ with $\lceil m \rceil x + \lceil b \rceil$ and $\lfloor m \rfloor x + \lceil b \rceil$ with $\lceil m \rceil x + \lceil b - \mathbf{err}(m)\underline{x} \rceil$.

Theorem 4 (Tightness of Class 1 Safe Linear Overestimators when $0 \notin [\underline{x}, \overline{x}]$). *Let g be a univariate function g and $mx + b$ be a linear overestimator for g in $[\underline{x}, \overline{x}]$. When $\underline{x} \geq 0$, $\lceil m \rceil x + b$ is tighter than $\lfloor m \rfloor x + \lceil b + \mathbf{err}(m)\overline{x} \rceil$. When $\overline{x} \leq 0$, $\lfloor m \rfloor x + \lceil b \rceil$ is tighter than $\lceil m \rceil x + \lceil b - \mathbf{err}(m)\underline{x} \rceil$.*

Proof. Consider the slightly tighter and easily comparable estimators $\lceil m \rceil x + b$ and $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$. The error of $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$ is:

$$\begin{aligned} E_1 &= \int_{\underline{x}}^{\overline{x}} \lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x} - g(x) dx \\ &= \frac{1}{2}(\overline{x} - \underline{x}) [\overline{x} (2 \lfloor m \rfloor - \lfloor m \rfloor - m) + \underline{x} (\lfloor m \rfloor - m)] + E \end{aligned}$$

where E is the error in $mx + b$. Likewise the error of $\lceil m \rceil x + b$ is:

$$\begin{aligned} E_2 &= \int_{\underline{x}}^{\overline{x}} \lceil m \rceil x + b - g(x) dx \\ &= \frac{1}{2}(\overline{x} - \underline{x}) (\lceil m \rceil - m)(\underline{x} + \overline{x}) + E \end{aligned}$$

$\lceil m \rceil x + b$ is tighter than $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$ when $E_2 < E_1$ which reduces to $\underline{x} < \overline{x}$. Since this condition is always met, $\lceil m \rceil x + b$ is always tighter than $\lfloor m \rfloor x + b + \mathbf{err}(m)\overline{x}$. Similarly, $\lfloor m \rfloor x + \lceil b \rceil$ is tighter than $\lceil m \rceil x + \lceil b - \mathbf{err}(m)\underline{x} \rceil$ when $\overline{x} \leq 0$. $\square$

Theorems 3 and 4 generalize and provide the theoretical justification for the heuristic used by Michel et al. [9].

4.2 Theoretical Results on Class 2 Estimators

We now study the tightness of Class-2 estimators. The next theorem compares the “real” counterparts of the two class-2 operators.

Theorem 5 (Tightness of Class 2 Safe Linear Overestimators). *Let g be a univariate function with linear overestimator $mx + b$ over $[\underline{x}, \bar{x}]$ such that $g(\underline{x}) = m\underline{x} + b$, $g(\bar{x}) = m\bar{x} + b$. $\lfloor m \rfloor x + g(\bar{x}) - \lfloor m \rfloor \bar{x}$ is a tighter estimator than $\lceil m \rceil x + g(\underline{x}) - \lceil m \rceil \underline{x}$ when $m - \lfloor m \rfloor < \lceil m \rceil - m$.*

Proof. The error of the $g(\bar{x}) = m\bar{x} + b$. $\lfloor m \rfloor x + g(\bar{x}) - \lfloor m \rfloor \bar{x}$ is given by

$$\begin{aligned} E_1 &= \int_{\underline{x}}^{\bar{x}} \lfloor m \rfloor x + g(\bar{x}) - \lfloor m \rfloor \bar{x} - g(x) \\ &= \frac{1}{2}(\bar{x} - \underline{x})^2(m - \lfloor m \rfloor) + E \end{aligned}$$

where E is the error of $mx + b$. Likewise the error in $\lceil m \rceil x + g(\underline{x}) - \lceil m \rceil \underline{x}$ is

$$E_2 = \frac{1}{2}(\bar{x} - \underline{x})^2(\lceil m \rceil - m) + E.$$

The $\lfloor m \rfloor$ formulation is tighter when $E_1 < E_2$ which reduces to $m - \lfloor m \rfloor < \lceil m \rceil - m$, i.e., when $\lfloor m \rfloor$ is a better approximation of m than $\lceil m \rceil$. $\square$

Of course, this result is not useful in practice, since m is not known and there is no way to evaluate the condition stated in Theorem 5. The theorem only considers the “real” counterparts of the estimators, i.e., it ignores the rounding errors in the actual evaluation of the operators. In other words, since these operators can be rewritten as $\lceil \lfloor m \rfloor (x - \bar{x}) + g(\bar{x}) \rceil$ and $\lceil \lceil m \rceil (x - \underline{x}) + g(\underline{x}) \rceil$, Theorem 5 applies to the Class-2 estimators whenever the rounding errors in these two terms are similar which in turn requires that $g(\underline{x}) \sim g(\bar{x})$. Fortunately, the next section, which relates both classes, gives a criterion to choose between them.

4.3 Theoretical Results on Class 1 and Class 2 Estimators

This section compares the Class 1 and Class 2 overestimators and describes the main result of this paper. The theorem states that Class-2 estimators are tighter than Class-1 estimators (modulo rounding errors). The theorem is given for the $\lceil m \rceil$ estimators, but a similar theorem can be proven for the $\lfloor m \rfloor$ estimators.

Theorem 6 (Relative Tightness of Class 1 and Class 2 Safe Linear Overestimators). *Let g be a univariate function with linear overestimator $mx + b$ over $[\underline{x}, \bar{x}]$ such that $g(\underline{x}) = m\underline{x} + b$. The class-2 estimators are tighter than the class-1 estimators (modulo rounding errors).*

Proof. We prove the result modulo rounding errors. When $\underline{x} \leq 0$, we have

$$\begin{aligned} &\lceil m \rceil x + \lceil g(\underline{x}) \rceil - \lceil m \rceil \underline{x} \\ &\leq \lceil m \rceil x + \lceil g(\underline{x}) - (m + \text{err}(m))\underline{x} \rceil \\ &\leq \lceil m \rceil x + \lceil b - \text{err}(m)\underline{x} \rceil \end{aligned}$$

which is the class-1 estimator when $\underline{x} \leq 0$. When $\underline{x} \geq 0$, we have

$$\begin{aligned} & \lceil m \rceil x + \lceil g(\underline{x}) - \lceil m \rceil \underline{x} \rceil \\ & \leq \lceil m \rceil x + \lceil g(\underline{x}) - m \underline{x} \rceil \\ & \leq \lceil m \rceil x + \lceil b \rceil \end{aligned}$$

which is the class-1 estimator when $\underline{x} \geq 0$. $\square$

Theorem 6 is interesting for many reasons. First, although it abstracts the rounding errors, it should hold in practice since the rounding errors in $g(\underline{x})$ should be smaller than, or similar to, those in b . (There are of course other terms but the errors cannot be radically different in the remaining parts). The experimental results in Section 4.4 confirms this. Second, it provides intuition as to why class-2 estimators are tighter than class-1 estimators. A class-1 operator systematically accounts for an error $\mathbf{err}(m) = \lceil m \rceil - \lfloor m \rfloor$ in the slope, while a class-2 estimator only adds an error $\lceil m \rceil - m$ (resp. $m - \lfloor m \rfloor$). Third, since class-2 operators can be viewed as tighter versions of the class-1 operators, similar criteria can be applied to choose between them, solving the problem left open in the previous section.

4.4 Numerical Results

We now compare numerically class-1 and class-2 estimators to confirm the findings of Theorem 6. More precisely, we compare $\lfloor m \rfloor x + \lceil g(\bar{x}) - \lfloor m \rfloor \bar{x} \rceil$ with $\lfloor m \rfloor x + \lceil b + \mathbf{err}(m)\bar{x} \rceil$ and $\lceil m \rceil x + \lceil g(\underline{x}) - \lceil m \rceil \underline{x} \rceil$ with $\lceil m \rceil x + \lceil b - \mathbf{err}(m)\underline{x} \rceil$ numerically for $0 \in [\underline{x}, \bar{x}]$ using the above heuristics to choose between the pairs. We use even powers for comparison purposes for which linear overestimators are given as follows.

$$g(x) = x^n \leq \underbrace{\frac{\bar{x}^n - \underline{x}^n}{\bar{x} - \underline{x}}}_m x + \underbrace{\frac{\bar{x}\bar{x}^n - \underline{x}\underline{x}^n}{\bar{x} - \underline{x}}}_b, \quad n \text{ even}, \quad x \in [\underline{x}, \bar{x}] \quad (3)$$

This general term can be simplified using $m = \underline{x} + \bar{x}$ and $b = -\underline{x}\bar{x}$ when $n = 2$. Moreover, since (3) is a secant of x^n , $g(x) = mx + b$ at both $x = \underline{x}$ and $x = \bar{x}$.

Given the theoretical results, it is easy to derive a set of numerical experiments. When $|\underline{x}| \geq |\bar{x}|$, we only compare the intercepts $\lceil g(\bar{x}) - \lfloor m \rfloor \bar{x} \rceil$ and $\lceil b + \mathbf{err}(m)\bar{x} \rceil$, using the estimators with the same slopes. The smallest intercept gives the tightest estimator. Likewise when $|\underline{x}| < |\bar{x}|$, we compare the intercepts $\lceil g(\underline{x}) - \lceil m \rceil \underline{x} \rceil$ and $\lceil b - \mathbf{err}(m)\underline{x} \rceil$.

Figures 5 and 6 depict the experimental results. To compute b , our numerical results use the specialized form for $n = 2$ (Figure 5) and the general form (Figure 6) otherwise. Random values were generated for $\underline{x}$ and $\bar{x}$ in a wide range of values. The results were collected region by region. We computed the percentage of cases where class-2 estimators were strictly tighter than class-1 estimators (%C21) and vice-versa (%C12). The figures report the difference (%C21 - %C12). For the quadratic case, Figure 5 shows that class-2 estimators




Fig. 5. Numerical Results for x^2 .

are very often tighter than class-1 operators. Typically, class-2 estimators are tighter in about 40-50% of the cases, while class-1 estimators are tighter in about 0-10% of the cases (they have equal tightness in the remaining cases). It is only when the two bounds are about the same size that class-1 improves over class-2 in about 40-50% of the cases. Figure 6 depicts the results for n^4 to n^{10} . The improvements of class-2 estimators here is striking. Class-2 estimators improve over class-1 estimators in more than 99% of the cases. Of course, this is not surprising in light of the proof of Theorem 6 since the slope is much steeper and hence the benefits of class-2 operators become more pronounced. Note also the improvement in tightness is proportional to the magnitude of the bounds. Overall, the experimental results confirm the theoretical results and clearly demonstrate the value of class-2 estimators.

5 Safe Linear Overestimators for Multivariate Functions

We now turn our attention to safe linear overestimators for multivariate functions. Multivariate functions frequently appear in global optimization. They can be estimated using a hyperplane in n dimensions. For example, a linear overestimator for the bilinear term xy (e.g., [5, 11]) is given by two planes:

$$xy \leq \min\{\underline{y}x + \bar{x}\bar{y} - \bar{x}\underline{y}, \bar{y}x + \underline{x}\underline{y} - \underline{x}\bar{y}\}, (x, y) \in [\underline{x}, \bar{x}] \times [\underline{y}, \bar{y}] \quad (4)$$

Since slopes of the two planes given in (4) are floating-point numbers, safe estimators for this case are given simply by rounding up the intercepts $-\bar{x}\bar{y}$ and $-\underline{x}\underline{y}$. The overestimator for the term $\frac{x}{y}$ used by [11, 16] have slopes that are




Fig. 6. Numerical Results for x^4 to x^{10} .

nonlinear functions of floating point numbers:

$$\frac{x}{y} \leq \min \left\{ \frac{\bar{y}}{\underline{y}^2} x + \frac{x}{\underline{y}^2} y + \frac{x\bar{y}}{\underline{y}^2}, \frac{y}{\bar{y}^2} x + \frac{\bar{x}}{\bar{y}^2} y + \frac{\bar{x}y}{\bar{y}^2} \right\}, (x, y) \in [\underline{x}, \bar{x}] \times [\underline{y}, \bar{y}], \underline{y} > 0 \quad (5)$$

In order to represent (5) with floating point numbers, special care must be taken in order to satisfy the two dimensional version of (1). More generally, estimators for multivariate functions can be obtained through estimators for factorable functions (see, for instance, [8]).

The above discussion indicates that it is critical in practice to generalize our results to multivariate functions. The problem can be formalized as follows (for overestimators): given an n -dimensional hyperplane overestimating an n -variate function $g(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$, $\mathbf{x} = (x_1, \dots, x_n)$, over $[\underline{x}_1, \bar{x}_1] \times \dots \times [\underline{x}_n, \bar{x}_n]$, the goal is to find $m_1^*, \dots, m_n^*, b^* \in \mathcal{F}$ such that:

$$\sum_{i=1}^n m_i^* x_i + b^* \geq \sum_{i=1}^n m_i x_i + b \geq g(\mathbf{x}), \quad \forall \mathbf{x} \in [\underline{x}_1, \bar{x}_1] \times \dots \times [\underline{x}_n, \bar{x}_n] \quad (6)$$

The first key result in this section is to show that safe linear estimators for multivariate functions can be derived naturally by combining univariate linear

estimators. In other words, the result makes it possible to consider each variable independently in the estimator

$$\sum_{i=1}^n m_i x_i + b,$$

replace m_i and b by their safe counterparts m_i^* and b_i^* , and combine all the individual coefficients into a safe estimator for the multivariate function. One of the interesting aspects of this result is the ability to factor b out from the b_i^* s to obtain a tight estimator.

Theorem 7 (Safe Linear Overestimators for Multivariate Functions). *Let g be an n -variate function and let $\sum_{i=1}^n m_i x_i + b$ be an overestimator for g in $[\underline{x}_1, \bar{x}_1] \times \cdots \times [\underline{x}_n, \bar{x}_n]$. Let $m_i^* x_i + b_i^*$ be a safe linear overestimator of $m_i x_i + b$ in $[\underline{x}_i, \bar{x}_i]$ and $\delta_i^* = \lceil b_i^* - b \rceil$, $1 \leq i \leq n$. Then, the hyperplane*

$$\sum_{i=1}^n m_i^* x_i + \lceil b + \sum_{i=1}^n \delta_i^* \rceil$$

is a safe linear overestimator for g in $[\underline{x}_1, \bar{x}_1] \times \cdots \times [\underline{x}_n, \bar{x}_n]$.

Proof. We show that

$$\sum_{i=1}^n m_i^* x_i + \lceil b + \sum_{i=1}^n \delta_i^* \rceil \geq \sum_{i=1}^n m_i x_i + b.$$

We have

$$\begin{aligned} \sum_{i=1}^n m_i^* x_i + \lceil b + \sum_{i=1}^n \delta_i^* \rceil &\geq \sum_{i=1}^n m_i^* x_i + b + \sum_{i=1}^n \delta_i^* \\ &\geq \sum_{i=1}^n (m_i^* x_i + b_i^* - b) + b \\ &= \sum_{i=1}^n (m_i^* x_i + b_i^*) - (n-1)b \\ &\geq \sum_{i=1}^n (m_i x_i + b) - (n-1)b \\ &= \sum_{i=1}^n m_i x_i + b. \quad \square \end{aligned}$$

Note that class-1 estimators trivially satisfy the hypotheses of this theorem. As a consequence, the theorem gives an elegant and simple way to obtain safe overestimators for multivariate functions. (A similar result can be derived for underestimators).

We now investigate the tightness of multivariate estimators derived using Theorem 7. For simplicity, we restrict attention to multivariate estimators derived by combining class-1 univariate estimators, which we call class-1 multivariate estimators. Our first result shows that class-1 multivariate estimators derived using Theorem 7 are as tight as possible for a given choice of rounding direction for each dimension (i.e., $\lfloor m_i \rfloor$ or $\lceil m_i \rceil$).

Theorem 8 (Tightness of Class-1 Safe Multivariate Overestimators). *Class-1 multivariate overestimators derived using Theorem 7 are as tight as possible for a given choice of rounding direction for each variable (i.e., $\lfloor m_i \rfloor$ or $\lceil m_i \rceil$).*

Proof. The safe overestimating hyperplane given by Theorem 7 can be written as:

$$\sum_{i \in I \cup K} \lceil m_i \rceil x_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor x_j + \lceil b - \sum_{k \in K} \text{err}(m_k) \underline{x}_k + \sum_{l \in L} \text{err}(m_l) \overline{x}_l \rceil$$

where sets I, J, K and L partition $\{1, \dots, n\}$. Using the same choices for $\lceil m \rceil$ and $\lfloor m \rfloor$ defined by this partition, we now calculate b^* such that $\sum_{i \in I \cup K} \lceil m_i \rceil x_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor x_j + b^* \geq \sum_{i=1}^n m_i x_i + b$. It is sufficient to satisfy this inequality at $(\tilde{x}_1, \dots, \tilde{x}_n)$ where

$$\tilde{x}_i = \begin{cases} \overline{x}_i & : i \in J \cup L \\ \underline{x}_i & : \text{otherwise} \end{cases}$$

due to the combination of $\lfloor m \rfloor$ and $\lceil m \rceil$ given by the partition. b^* must satisfy

$$\sum_{i \in I \cup K} \lceil m_i \rceil \underline{x}_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor \overline{x}_j + b^* \geq \sum_{i \in I \cup K} m_i \underline{x}_i + \sum_{j \in J \cup L} m_j \overline{x}_j + b.$$

By a case analysis on the sign of $\tilde{x}_i$ for each i :

$$\begin{aligned} & \sum_{i \in I \cup K} m_i \underline{x}_i + \sum_{j \in J \cup L} m_j \overline{x}_j + b \\ & \leq \sum_{i \in I} \lceil m_i \rceil \underline{x}_i + \sum_{j \in J} \lfloor m_j \rfloor \overline{x}_j + \sum_{k \in K} \lfloor m_k \rfloor \underline{x}_k + \sum_{l \in L} \lceil m_l \rceil \overline{x}_l + b \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{i \in I \cup K} \lceil m_i \rceil \underline{x}_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor \overline{x}_j + b^* \\ & \geq \sum_{i \in I} \lceil m_i \rceil \underline{x}_i + \sum_{j \in J} \lfloor m_j \rfloor \overline{x}_j + \sum_{k \in K} \lfloor m_k \rfloor \underline{x}_k + \sum_{l \in L} \lceil m_l \rceil \overline{x}_l + b. \end{aligned}$$

Collecting terms: $b^* \geq b - \sum_{k \in K} \text{err}(m_k) \underline{x}_k + \sum_{l \in L} \text{err}(m_l) \overline{x}_l$. Correctly rounding this results in the same hyperplane as constructed by the theorem. $\square$

It remains to choose appropriate rounding directions for each variable. The next theorem shows that the combination of tight class-1 univariate estimators gives a tight class-1 multivariate estimator.

Theorem 9 (Optimality of Class-1 Safe Multivariate Overestimators).
A multivariate class-1 safe estimators derived using Theorem 7 by choosing optimal class-1 univariate safe estimators for each variable is an optimal class-1 multivariate safe estimator.

Proof. Consider the hyperplane:

$$H = \sum_{i \in I \cup K} \lceil m_i \rceil x_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor x_j + \lceil b - \sum_{k \in K} \text{err}(m_k) \underline{x}_k + \sum_{l \in L} \text{err}(m_l) \bar{x}_l \rceil \quad (7)$$

where $I = \{i : \underline{x}_i \geq 0\}$, $J = \{j : \bar{x}_j \leq 0\}$, $K = \{k : 0 \in (\underline{x}_k, \bar{x}_k), |\underline{x}_k| < |\bar{x}_k|\}$ and $L = \{l : 0 \in (\underline{x}_l, \bar{x}_l), |\underline{x}_l| \geq |\bar{x}_l|\}$. By theorem 7, H is safe. By theorem 8, H is as tight as possible given the choices of $\lceil m \rceil$ and $\lfloor m \rfloor$ defined by the partition. Consider the error between a slightly tighter version of (7), given by an unrounded intercept, and $\sum_{i=1}^n m_i x_i + b$. The remaining error $(\int_{\underline{x}_1}^{\bar{x}_1} dx_1 \cdots \int_{\underline{x}_n}^{\bar{x}_n} dx_n \sum_{i=1}^n m_i x_i + b - g)$ is constant over all 2^n possible safe class-1 overestimating hyperplanes. The error is defined by the natural extension of definition 3 to higher dimensions:

$$\begin{aligned} E &= \int_{\underline{x}_1}^{\bar{x}_1} dx_1 \cdots \int_{\underline{x}_n}^{\bar{x}_n} dx_n \left\{ \sum_{i \in I \cup K} \lceil m_i \rceil x_i + \sum_{j \in J \cup L} \lfloor m_j \rfloor x_j + b - \sum_{k \in K} \text{err}(m_k) \underline{x}_k \right. \\ &\quad \left. + \sum_{l \in L} \text{err}(m_l) \bar{x}_l - \left(\sum_{i=1}^n m_i x_i + b \right) \right\} \\ &= \int_{\underline{x}_1}^{\bar{x}_1} dx_1 \cdots \int_{\underline{x}_n}^{\bar{x}_n} dx_n \left\{ \sum_{i \in I \cup K} (\lceil m_i \rceil - m_i) x_i + \sum_{j \in J \cup L} (\lfloor m_j \rfloor - m_j) x_j \right. \\ &\quad \left. - \sum_{k \in K} \text{err}(m_k) \underline{x}_k + \sum_{l \in L} \text{err}(m_l) \bar{x}_l \right\} \end{aligned}$$

Using the integrals

$$\begin{aligned} \int_{\underline{x}_1}^{\bar{x}_1} dx_1 \cdots \int_{\underline{x}_n}^{\bar{x}_n} dx_n 1 &= \prod_{j=1}^n (\bar{x}_j - \underline{x}_j) = P \\ \int_{\underline{x}_1}^{\bar{x}_1} dx_1 \cdots \int_{\underline{x}_n}^{\bar{x}_n} dx_n x_i &= \frac{1}{2} (\bar{x}_i^2 - \underline{x}_i^2) \prod_{j \neq i} (\bar{x}_j - \underline{x}_j) \\ &= \frac{1}{2} (\bar{x}_i + \underline{x}_i) P \end{aligned}$$

the error is rewritten as:

$$\begin{aligned}
E &= P \left\{ \sum_{i \in I \cup K} \frac{1}{2} (\lceil m_i \rceil - m_i) (\bar{x}_i + \underline{x}_i) + \sum_{j \in J \cup L} \frac{1}{2} (\lfloor m_j \rfloor - m_j) (\bar{x}_i + \underline{x}_i) \right. \\
&\quad \left. - \sum_{k \in K} \text{err}(m_k) \underline{x}_k + \sum_{l \in L} \text{err}(m_l) \bar{x}_l \right\} \\
&= \frac{P}{2} \left\{ \sum_{i \in I} (\lceil m_i \rceil - m_i) (\bar{x}_i + \underline{x}_i) + \sum_{j \in J} (\lfloor m_j \rfloor - m_j) (\bar{x}_i + \underline{x}_i) \right. \\
&\quad + \sum_{k \in K} \{ (\lceil m_i \rceil - m_i) \bar{x}_i + (2 \lfloor m_i \rfloor - \lceil m_i \rceil - m_i) \underline{x}_i \} \\
&\quad \left. + \sum_{l \in L} \{ (\lfloor m_i \rfloor - m_i) \underline{x}_i + (2 \lceil m_i \rceil - \lfloor m_i \rfloor - m_i) \bar{x}_i \} \right\} \tag{8}
\end{aligned}$$

Define the type of a variable as the set (one of I , J , K , or L) to which it belongs. We show that another partition, $\{I', J', K', L'\}$, does not define a tighter hyperplane, H' . Since the error given by (8) is linear in the type of variable, we can consider each variable independently. H and H' can differ in any of the following four ways while remaining safe:

- A variable, x_i , in set I can be moved to set L producing $I' = I \setminus \{x_i\}$ and $L' = L \cup \{x_i\}$. This will add the term $P \cdot \text{err}(m_l) \bar{x}_l \geq 0$ to the error, thereby increasing the total error. H remains tighter than H' .
- A variable, x_i , in set J can be moved to set K producing $J' = J \setminus \{x_i\}$ and $K' = K \cup \{x_i\}$. This will add the term $-P \cdot \text{err}(m_l) \underline{x}_l \geq 0$ to the error, thereby increasing the total error. H remains tighter than H' .
- A variable, x_i in set K can move to set L producing $K' = K \setminus \{x_i\}$ and $L' = L \cup \{x_i\}$. The difference in error is:

$$\begin{aligned}
&\frac{P}{2} \{ (\lfloor m_i \rfloor - m_i) \underline{x}_i + (2 \lceil m_i \rceil - \lfloor m_i \rfloor - m_i) \bar{x}_i \\
&\quad - (\lceil m_i \rceil - m_i) \bar{x}_i - (2 \lfloor m_i \rfloor - \lceil m_i \rceil - m_i) \underline{x}_i \} \\
&= P \cdot \text{err}(m) \{ \bar{x}_i + \underline{x}_i \} \\
&\geq 0 \quad \text{since the variable in } K \text{ satisfies } |\bar{x}| > |\underline{x}|.
\end{aligned}$$

Since the error increases with this change, H remains tighter than H' .

- Likewise, a variable moving from set L to set K increases the error of the overestimating hyperplane.

By the above arguments, the hyperplane defined by theorem 7 created by combining tight safe class-1 univariate overestimators is the tightest class-1 overestimator for an n -variate function. $\square$

6 Conclusion

Global optimization problems are often approached by branch and bound algorithms which use linear relaxations of the nonlinear constraints computed from the current variable bounds at each node of the search tree. This paper considered the problem of obtaining safe linear relaxations which are guaranteed to enclose the solutions of the nonlinear problem. It made two main contributions. On the one hand, it studied two classes of linear estimators univariate functions. The first class of estimators generalizes the results in [9] from quadratic to arbitrary functions, while the second class is entirely novel. Theoretical and numerical results indicated that class-2 estimators, when they apply, are tighter than class-1 estimators. On the other hand, the paper generalized the univariate results to multivariate functions. It indicated how to derive estimators for multivariate functions by combining univariate estimators derived for each variable independently. Moreover, it showed that the combination of tight class-1 safe univariate estimators is a tight class-1 safe multivariate estimators. Together these results provide a systematic, comprehensive, and elegant framework to derive safe linear estimators for global optimization problems. In conjunction with the safe bounds on the linear relaxations derived in [10, 4], they provide the theoretical foundation for rigorous results of branch and bound approaches to global optimization based on linear programming. Future work aims at studying how to generalize the multivariate tightness results to class-2 estimators.

References

1. F. Benhamou, D. McAllester, and P. Van Hentenryck. CLP(Intervals) Revisited. In *Proceedings of the International Symposium on Logic Programming (ILPS-94)*, pages 124–138, Ithaca, NY, November 1994.
2. F. Benhamou and W. Older. Applying Interval Arithmetic to Real, Integer and Boolean Constraints. *Journal of Logic Programming*, 1997. To appear.
3. I. E. Grossmann and S. Lee. Generalized convex disjunctive programming: Non-linear convex hull relaxation. Submitted for publication (2001).
4. C. Jansson. Rigorous lower and upper bounds in linear programming, 2002.
5. Y. Lebbah, M. Rueher, and C. Michel. A global filtering algorithm for handling systems of quadratic equations and inequations. *Lecture Notes in Computer Science*, 2470:109–123, 2002.
6. S. Lee and I. E. Grossmann. A global optimization algorithm for nonconvex generalized disjunctive programming and applications to process systems. *Computers and Chemical Engineering*, 25:1675–1697, 2001.
7. O. Lhomme. Consistency Techniques for Numerical Constraint Satisfaction Problems. In *Proceedings of the 1993 International Joint Conference on Artificial Intelligence*, Chambery, France, August 1993.
8. G. P. McCormick. Computability of global solutions to factorable nonconvex programs: Part I - convex underestimating problems. *Mathematical Programming*, 10:146–175, 1976.
9. C. Michel, Y. Lebbah, and M. Rueher. Safe embedding of the simplex algorithm in a CSP framework. *Proceedings CPAIOR'03*, pages 210–220, 2003.

10. A. Neumaier and O. Shcherbina. Safe bounds in linear and mixed-integer programming. to appear in Mathematical Programming.
11. I. Quesada and I. E. Grossmann. A global optimization algorithm for linear fractional and bilinear programs. *Journal of Global Optimization*, 6:39–76, 1995.
12. H. S. Ryoo and N. V. Sahinidis. Global optimization of nonconvex NLPs and MINLPs with applications in process design. *Computer and Chemical Engineering*, 19:551–566, 1995.
13. P. Van Hentenryck, D. McAllister, and D. Kapur. Solving Polynomial Systems Using a Branch and Prune Approach. *SIAM Journal on Numerical Analysis*, 34(2), 1997.
14. P. Van Hentenryck, L. Michel, and Y. Deville. *Numerica: a Modeling Language for Global Optimization*. The MIT Press, Cambridge, Mass., 1997.
15. J. M. Zamora and I. E. Grossmann. A comprehensive global optimization approach for the synthesis of heat exchanger networks with no stream splits. *Computers and Chemical Engineering*, 21:S65–S70, 1997.
16. J. M. Zamora and I. E. Grossmann. A branch and contract algorithm for problems with concave univariate, bilinear and linear fractional terms. *Journal of Global Optimization*, 14:217–249, 1999.